

Recombinant Mouse IL-3 (C-6His)

Catalog No : PMK2140

Known As: Interleukin-3; IL-3; Hematopoietic growth factor; Multipotential colony-stimulating factor; P-cell-stimulating factor; Il3; Il-3; Mast cell growth factor; MCGF

PROPERTIES

Description	Recombinant Mouse Interleukin-3 is produced by our Mammalian expression system and the target gene encoding Ala27-Cys166 is expressed with a 6His tag at the Cterminus.
Accession	P01586
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Size	10μg/50μg/500μg/1mg
Purity	> 95%
Endotoxin	< 0.01 EU/μg as determined by LAL test.
Predicted Mol Mass	16.5 KDa
Apparent Mol Mass	15-32 KDa, reducing conditions
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Background	Interleukin 3 is a pleiotropic factor produced primarily by activated T cells that can stimulate the proliferation and differentiation of pluripotent hematopoietic stem cells as well as various lineage committed progenitors. In addition, IL-3 also affects the functional activity of mature mast cells, basophils, eosinophils and macrophages. Because of its multiple functions and targets, it was originally studied under different names, including mast cell growth factor P-cell stimulating factor, burst promoting activity, multi-colony stimulating factor, thy-1 inducing factor and WEHI-3 growth factor. In addition to activated T cells, other cell types such as human thymic epithelial cells, activated mouse mast cells, mouse keratinocytes and neurons/astrocytes can also produce IL-3. IL-3 exerts its biological activities through binding to specific cell surface receptors. The high affinity receptor responsible for IL- 3. signaling is composed of α and β subunits. IL-3 is capable of supporting the proliferation of a broad range of hematopoietic cell types. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. IL-3 has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders.

NOTE: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.