

## Recombinant Human IL-2

Catalog No : PMK2121

Known As: Interleukin-2; IL-2; T-Cell Growth Factor; TCGF; Aldesleukin; IL2

### PROPERTIES

Description	Recombinant Human Interleukin-2 is produced by our E.coli expression system and the target gene encoding Ala21-Thr153 is expressed.
Accession	P60568
Formulation	Lyophilized from a 0.2 µm filtered solution of 10mM sodium citrate, pH 4.0.
Size	10µg/50µg/500µg/1mg
Purity	> 95%
Endotoxin	< 0.01 EU/µg as determined by LAL test.
Predicted Mol Mass	15.5 KDa
Apparent Mol Mass	14 KDa, reducing conditions
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Background	Interleukin-2(IL-2) is an interleukin, a type of cytokine signaling molecule in the immune system, belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL- 2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells, NK cells, lymphokine-activated killer cells, monocytes, macrophages and oligodendrocytes.

**NOTE:** The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.